

Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books  
Search PubMed for Go Clear  
Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

☐ 1: Development. 1995 Jun;121(6):1787-99.

[Related Articles, Links](#)

Entrez PubMed

Overview  
Help | FAQ  
Tutorial  
New/Noteworthy  
E-Utilities

FREE full text article at  
[dev.biologists.org](http://dev.biologists.org)

## **Zebrafish wnt8 and wnt8b share a common activity but are involved in distinct developmental pathways.**

**Kelly GM, Greenstein P, Erezyilmaz DF, Moon RT.**

Howard Hughes Medical Institute, Department of Pharmacology, Seattle, Washington, USA.

The specification of the vertebrate body plan is dependent on numerous signaling molecules, including members of the Wnt family. We have identified two zebrafish wnt8 paralogs related to Xwnt-8B and Xwnt-8, respectively. A RT-PCR assay demonstrated that wnt8 is expressed maternally, with transcripts detected throughout embryogenesis, whereas wnt8b transcripts were first detected during late gastrulation. The wnt8 transcripts at 50% epiboly are spatially restricted to those cells at the blastoderm margin, overlying gsc-expressing cells in the axial hypoblast. During late gastrulation, wnt8 was no longer detected in the marginal cells at the dorsal midline and by mid-segmentation, transcripts were found in the presumptive tail bud. In contrast, wnt8b expression is spatially restricted to prospective neuroepithelium, and later to neural-specific structures. Overexpression of both wnts results in two major phenotypes: radialized embryos and embryos with anterior defects. These phenotypes were preceded by significant changes in the spatial expression patterns of gsc and ntl transcripts, reminiscent of activities of Xwnt-8 in *Xenopus*, and consistent with a role for wnt8 in the specification or patterning of mesoderm.


PMID: 7600994 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text

[Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Jul 27 2004 06:47:37

## BEST AVAILABLE COPY



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books  
Search PubMed for Go Clear  
Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

☐ 1: Mech Dev. 1998 Feb;71(1-2):119-29.[Related Articles, Links](#)

Entrez PubMed

Overview  
Help | FAQ  
Tutorial  
New/Noteworthy  
E-Utilities

PubMed Services

Journals Database  
MeSH Database  
Single Citation Matcher  
Batch Citation Matcher  
Clinical Queries  
LinkOut  
Cubby

Related Resources

Order Documents  
NLM Gateway  
TOXNET  
Consumer Health  
Clinical Alerts  
ClinicalTrials.gov  
PubMed Central

## BMP-2/-4 and Wnt-8 cooperatively pattern the Xenopus mesoderm.

Hoppler S, Moon RT.

Howard Hughes Medical Institute and Department of Pharmacology, University of Washington School of Medicine, Seattle 98195, USA.

Establishment of the dorsoventral axis is central to animal embryonic organization. In *Xenopus* two different classes of signaling molecules function in the dorsoventral patterning of the mesoderm. Both the TGF-beta-related products of the BMP-2 and BMP-4 genes and the Wnt molecule encoded by *Xenopus* Wnt-8 specify ventral fate and appear to inhibit dorsal mesodermal development. The similar functions of these molecularly very different classes of signaling molecules prompted us to study their mutual regulation and to closely compare their roles in mesoderm patterning. We find that Wnt-8 and BMP-4 are indistinguishable in their abilities to induce expression of ventral genes. Although BMP-2/-4 signaling regulates Wnt-8 expression, these genes do not function in a linear pathway because Wnt-8 overexpression cannot compensate for an inhibition of BMP-2/-4 function, but rather BMP-4 overexpression rescues ventral gene expression in embryos with inhibited Wnt-8 function. We further find that Wnt-8 and BMP-2/-4 differ in their abilities to regulate dorsal gene expression. While BMP-4 appears to generally inhibit the expression of dorsal genes, *Xenopus* Wnt-8 only inhibits the expression of the notochord marker *Xnot*. Whereas the inhibitory effect of BMP-2/-4 localizes dorsal mesodermal fate, our results suggest that *Xenopus* Wnt-8 functions in the further patterning of the dorsal mesoderm into the most dorsal sector from which the notochord develops and the dorsolateral sector from where the somites differentiate.

PMID: 9507084 [PubMed - indexed for MEDLINE]


Display Abstract Show: 20 Sort Send to Text

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)  
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Jul 27 2004 06:47:37

BEST AVAILABLE COPY



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for [ ] Go Clear

Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

☐ 1: Development. 1998 Nov;125(21):4283-92.[Related Articles, Links](#)

Entrez PubMed

Overview  
Help | FAQ  
Tutorial  
New/Noteworthy  
E-Utilities

FREE full-text article at  
[dev.biologists.org](http://dev.biologists.org)

## Evidence for a frizzled-mediated wnt pathway required for zebrafish dorsal mesoderm formation.

Nasevicius A, Hyatt T, Kim H, Guttman J, Walsh E, Sumanas S, Wang Y, Ekker SC.

Department of Biochemistry and Institute of Human Genetics, University of Minnesota Medical School, Minneapolis, MN 55455, USA.

We have used zebrafish as a model system for the study of vertebrate dorsoventral patterning. We isolated a maternally expressed and dorsal organizer localized member of the frizzled family of wnt receptors. Wild-type and dominant, loss-of-function molecules in misexpression studies demonstrate frizzled function is necessary and sufficient for dorsal mesoderm specification. frizzled activity is antagonized by the action of GSK-3, and we show GSK-3 is also required for zebrafish dorsal mesoderm formation. frizzled cooperatively interacts with the maternally encoded zebrafish wnt8 protein in dorsal mesodermal fate determination. This frizzled-mediated wnt pathway for dorsal mesoderm specification provides the first evidence for the requirement of a wnt-like signal in vertebrate axis determination.

PMID: 9753682 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text


[Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)

Department of Health &amp; Human Services

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Jul 27 2004 06:47:37

BEST AVAILABLE COPY



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for Go Clear

Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

1: Genes Dev. 1999 Dec 1;13(23):3149-59.

Related Articles, Links

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

FREE full text article at  
[www.genesdev.org](http://www.genesdev.org)

## Wnt signaling in *Xenopus* embryos inhibits bmp4 expression and activates neural development.

Baker JC, Beddington RS, Harland RM.

Department of Molecular and Cell Biology, University of California, Berkeley, California 94720, USA.

We report a new role for Wnt signaling in the vertebrate embryo: the induction of neural tissue from ectoderm. Early expression of mouse *wnt8*, *Xwnt8*, *beta*-catenin, or dominant-negative GSK3 induces the expression of neural-specific markers and inhibits the expression of *Bmp4* in *Xenopus* ectoderm. We show that *Wnt8*, but not the BMP antagonist *Noggin*, can inhibit *Bmp4* expression at early gastrula stages. Furthermore, inhibition of *beta*-catenin activity in the neural ectoderm of whole embryos by a truncated TCF results in a decrease in neural development. Therefore, we suggest that a cleavage-stage Wnt signal normally contributes to an early repression of *Bmp4* on the dorsal side of the embryo and sensitizes the ectoderm to respond to neural inducing signals from the organizer. The Wnt targets *Xnr3* and *siamois* have been shown previously to have neuralizing activity when overexpressed. However, antagonists of Wnt signaling, *dnXwnt8* and *Nxfzr8*, inhibit Wnt-mediated *Xnr3* and *siamois* induction, but not neural induction, suggesting an alternative mechanism for *Bmp* repression and neuralization. Conversely, *dnTCF* blocks both Wnt-mediated *Xnr3* and neural induction, suggesting that both pathways require this transcription factor.

PMID: 10601040 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Jul 27 2004 06:47:37